



## FACT SHEET No. 6

# Palliative Care for the Older Person in Pain

Worldwide, the population over the age of 60 years will double by 2050 [11]. As people live longer, increasing numbers will live with and die from multi-morbidity, frailty, and chronic health conditions such as renal or cardiac failure. In addition, older people may experience significant psychosocial stressors such as bereavement and loss of independence.

### What is palliative care?

Palliative care aims to maintain or improve quality of life and alleviate suffering, through early identification, detailed assessment, and treatment of symptoms [4]. In older people this:

- Combines geriatric medicine and palliative care focusing on comprehensive assessment to integrate social, spiritual, and environmental factors.
- Requires an understanding of multi-morbidity, safe prescribing, and a multidisciplinary approach.
- Prioritizes good communication, considering autonomy, involvement in decision-making, and the existence of ethical dilemmas.
- Works with older people and their families across settings (home, long-term care, hospices, and hospital) and during transitions [7].

### Pain assessment

How older people experience and report pain is mediated by a range of social and psychological factors, including stoicism, which may lead to under-reporting of pain [2]. The 'gold standard' remains selfreport. [3]. Questions about pain include three key dimensions: 1) sensory, 2) affective and 3) impact [8].

### Dementia and cognitive impairment

Reporting pain can be challenging for older people with cognitive impairment secondary to dementia and other neurodegenerative diseases, strokes, cultural, or language factors. Many people with dementia can



© Copyright 2019 International Association for the Study of Pain. All rights reserved.

**IASP brings together scientists, clinicians, health-care providers, and policymakers to stimulate and support the study of pain and translate that knowledge into improved pain relief worldwide.**



© Copyright 2019 European Pain Federation EFIC. All rights reserved.

**The European Pain Federation EFIC is a multidisciplinary professional organization in the field of pain research and medicine, consisting of the 37 European IASP Chapters**

report pain reliably [12], but it is essential to obtain collateral history. Direct observation or validated observational pain scales recognize how pain or discomfort may lead to behavioral change [9]. The American Geriatrics Society Guidelines [3] include a range of indicators:

	<b>Domain</b>	<b>Example</b>
1	Facial expressions	Frowning
2	Verbalizations and vocalizations	Moaning, grunting
3	Body movements	Guarding an area of the body, pacing
4	Changes in interpersonal interactions	Becoming withdrawn, aggression
5	Changes in activity patterns or routines	Appetite, activities of daily living, sleep
6	Mental status changes	Delirium, tearfulness, crying

Most observational pain tools contain items from these domains. Commonly used tools include the Abbey Pain Scale [1], Pain Assessment in Advanced Dementia (PAINAD) [10], and the Pain Assessment Checklist for Seniors with Limited Ability to Communicate (PACSLAC) [5].

### **Principles of management**

Non-pharmacological management, such as exercise, assistive devices or relaxation [2], or topical preparations, including NSAIDS for localized musculoskeletal pain [2], may be effective as a first choice. Pharmacological treatment of pain in older people may be challenging. Polypharmacy is common and changes in how drugs are metabolized and excreted increase the risk of interactions and side effects. Recommendations have been made [2] (AGS 2009) to lessen the risk of adverse events:

- Prescribe using the WHO pain ladder (<http://www.who.int/cancer/palliative/painladder/en/>).
- Start low with one drug and increase the dose slowly to achieve response. •
- Use the least invasive route of administration.



© Copyright 2019 International Association for the Study of Pain. All rights reserved.

**IASP brings together scientists, clinicians, health-care providers, and policymakers to stimulate and support the study of pain and translate that knowledge into improved pain relief worldwide.**



© Copyright 2019 European Pain Federation EFIC. All rights reserved.

**The European Pain Federation EFIC is a multidisciplinary professional organization in the field of pain research and medicine, consisting of the 37 European IASP Chapters**

	<b>Indication</b>	<b>Notes</b>
Paracetamol/ acetaminophen	Musculoskeletal pain and osteoarthritis	Effective and safe analgesic
Non-steroidal anti-inflammatory drugs i.e. naproxen, ibuprofen	Musculoskeletal pain and osteoarthritis where paracetamol not effective	Higher risk of adverse events. GI bleeding, cardiovascular side effects (increased risk of arterial hypertension, heart failure) and worsening of chronic kidney disease.
Codeine	Weak opioid for moderate pain	Sedation, hallucinations, delirium, nausea, vomiting, constipation, urinary retention, falls, fractures. Fentanyl or buprenorphine available as patches but cannot be used in opioid naïve patients
Morphine, oxycodone, fentanyl	Strong opioids for severe pain in cancer and non-cancer pain	
Amitriptyline	Neuropathic pain	Postural hypotension, cardiac arrhythmias, urinary retention, glaucoma and worsening cognitive functioning
Pregabalin, gabapentin	Neuropathic pain	Anxiolytic and sedative effect

## Conclusion

A palliative care approach ensures symptoms and goals of treatment are regularly reviewed. Discussing a “ceiling of care” with the person or their family to develop a treatment escalation plan supports good pain management by considering decisions such as discontinuing painful interventions. It may reduce the risks of people undergoing distressing transfers towards the end of life, particularly into hospitals [6]. In palliative care, we consider the concept of ‘total pain’ exploring psychological distress, which may influence the perception of pain and suffering.

## REFERENCES

[1] Abbey J, Piller N, De BA, Esterman A, Parker D, Giles L, Lowcay B. The Abbey pain scale: a 1-minute numerical indicator for people with end-stage dementia. *IntJ Palliat Nurs* 2004;10(1):6-13.

[2] Abdulla A, Bone M, Adams N, Elliott AM, Jones D, Knaggs R, Martin D, Sampson EL, Schofield P. Evidence-based clinical practice guidelines on management of pain in older people. *Age Ageing* 2013;42(2):151-153.

[3] American Geriatrics Society. The management of persistent pain in older persons. *J Am Geriatr Soc* 2002;50(6 Suppl):S205-



© Copyright 2019 International Association for the Study of Pain. All rights reserved.

**IASP brings together scientists, clinicians, health-care providers, and policymakers to stimulate and support the study of pain and translate that knowledge into improved pain relief worldwide.**



© Copyright 2019 European Pain Federation EFIC. All rights reserved.

**The European Pain Federation EFIC is a multidisciplinary professional organization in the field of pain research and medicine, consisting of the 37 European IASP Chapters**

224.

[4] Davies E, Higginson IJ. Better palliative care for older people, 2004.

[5] Fuchs-Lacelle S, Hadjistavropoulos T. Development and preliminary validation of the pain assessment checklist for seniors with limited ability to communicate (PACSLAC). *Pain ManagNurs* 2004;5(1):37-49.

[6] Obolensky L, Clark T, Matthew G, Mercer M. A patient and relative centred evaluation of treatment escalation plans: a replacement for the do-not-resuscitate process. *J Med Ethics* 2010;36(9):518-520.

[7] Pautex S, Curiale V, Pfisterer M, Rexach L, Ribbe M, Van Den Noortgate N. A common definition of geriatric palliative medicine. *J Am Geriatr Soc* 2010;58(4):790-791.

[8] Royal College of Physicians, British Geriatrics Society, British Pain Society. The assessment of pain in older people: national guidelines. Concise guidance to good practice series, Vol. 8, 2007.

[9] Scherder E, Herr K, Pickering G, Gibson S, Benedetti F, Lautenbacher S. Pain in dementia. *Pain* 2009;145(3):276-278.

[10] Warden V, Hurley AC, Volicer L. Development and psychometric evaluation of the Pain Assessment in Advanced Dementia (PAINAD) scale. *JAmMedDirAssoc* 2003;4(1):9-15.

[11] World Health Organisation. Global Health Observatory (GHO) data; mortality and global health estimates, Vol. 2017, 2017.

[12] Zwakhalen SM, Hamers JP, Berger MP. The psychometric quality and clinical usefulness of three pain assessment tools for elderly people with dementia. *Pain* 2006;126(1-3):210-220.

## AUTHOR

Elizabeth L Sampson, PhD  
Marie Curie Palliative Care Research Department,  
University College London  
London, United Kingdom

Sophie Pautex, MD  
Division of Palliative Medicine  
University Hospital Geneva, Geneva University  
Geneva, Switzerland



© Copyright 2019 International Association for the Study of Pain. All rights reserved.

**IASP brings together scientists, clinicians, health-care providers, and policymakers to stimulate and support the study of pain and translate that knowledge into improved pain relief worldwide.**



© Copyright 2019 European Pain Federation EFIC. All rights reserved.

**The European Pain Federation EFIC is a multidisciplinary professional organization in the field of pain research and medicine, consisting of the 37 European IASP Chapters**

[Please be sure to translate the text inside this box and on the next page as well as the text in the footer below.]

#### **About the International Association for the Study of Pain®**

IASP is the leading professional forum for science, practice, and education in the field of pain. [Membership is open to all professionals](#) involved in research, diagnosis, or treatment of pain. IASP has more than 7,000 members in 133 countries, 90 national chapters, and 22 Special Interest Groups.

#### **About the European Pain Federation EFIC®**

The European Pain Federation EFIC is a multidisciplinary professional organization in the field of pain research and medicine, consisting of the 37 European Chapters of the International Association for the Study of Pain (IASP) and representing some 20,000 physicians, nurses, scientists, psychologists, physiotherapists and other health care professionals involved in pain medicine

**As part of the Global and European Year Against Pain in the Most Vulnerable, IASP and EFIC offers a series of Fact Sheets that cover specific topics related to pain. These documents have been translated into multiple languages and are available for free download. Visit [GYAP Page](#) and [EYAP Page](#) more information.**



© Copyright 2019 International Association for the Study of Pain. All rights reserved.

**IASP brings together scientists, clinicians, health-care providers, and policymakers to stimulate and support the study of pain and translate that knowledge into improved pain relief worldwide.**



© Copyright 2019 European Pain Federation EFIC. All rights reserved.

**The European Pain Federation EFIC is a multidisciplinary professional organization in the field of pain research and medicine, consisting of the 37 European IASP Chapters**