



# 2020 EUROPEAN YEAR FOR THE PREVENTION OF PAIN



## Factsheet

### Secondary and Tertiary Prevention of Chronic Pain

#### Introduction

Prevention of pain and its chronic forms will reduce the burden of suffering for individuals and society, respectively. Secondary prevention aims to detect disease in the early stages and limits its progression [1, 2], whereby tertiary prevention aims to reduce or avoid complications or limiting consequences of the disease already present [1]. For pain, the definitions state to prevent individuals from developing chronic pain after initial onset of acute pain (secondary prevention), or to reduce ongoing disability, disuse or loss of social contacts and occupation, once pain has become chronic (tertiary prevention). Even though the research body for secondary pain prevention has grown during the past years, the focus remains on treating chronic pain.

The Global Year for the Prevention of Pain is a chance to raise awareness on strategies and interventions for the primary and secondary stages of disease development, supporting sufferers in maintaining daily physical, personal and social activities the best way possible to reduce the development of chronic pain. Recommendations in designing prevention trials have been published [3], which will hopefully generate reliable evidence.

#### Secondary Prevention of Pain: Preventing Chronicity

The secondary prevention of pain was first described by Fordyce in the 1970s [4], when distinguishing between pain and pain behaviour (disabling behaviour). A prevention regime was applied to reduce pain behaviour, leading to preliminary promising results in terms of prevention strategies [2, 4] including time-contingent training and medication. Secondary prevention requires an understanding of the factors involved in chronification, validated instruments to detect patients at risk, and validated clinical strategies addressing these risk factors with specific interventions [5]. Several reviews report risk factors for maintaining different forms of pain; red (biological) and yellow (psychosocial) flags for musculoskeletal pain is the most common construct, followed by blue (occupational), black (compensation) and white (socio-cultural) [6] flags, where evidence is still pending regarding their impact in secondary prevention of pain. Specific mechanism-based risk factors would help designing preventive strategies providing future base of tailored interventions [5, 7]. Such risk factors have been identified in catastrophizing/maladaptive cognitions and depressive mood for developing chronic knee [8] or low back pain [9] as yellow flags, or joint damage as red flag for knee pain [8]. Others reported several risk factors for developing chronic pain after surgery (e.g. [10-12]) with sound evidence for age, gender, type of surgery, genetic factors, preceding pain or history of chronic pain of other origin and a variety of psychosocial factors. Screening instruments are available especially for back pain with

preliminary evidence for their prognostic value [13]. Preliminary evidence is also provided that subgrouping patients regarding their risk for chronification and specifically tailored treatment is effective in the short and middle terms [14]. Education and exercise in preventing back pain from becoming chronic seems to be of best evidence at the moment [5, 7], primarily in the short and middle terms.

In contrast, iatrogenic factors (factors lying in the health care system) seem to contribute to chronification of pain, especially when solely focusing on somatic factors, ignoring multi-causal genesis of pain, overestimating the impact of somatic or radiologic findings on wellbeing of the patient and overuse of diagnostic procedures and rather passively oriented interventions in long term (e.g. massage) [15]. In light of these findings, responsibility of health care providers in treating pain in an evidence-based manner should not be underestimated and should be included into research activities.

Intensified effort in establishing models [3] such as the avoidance-endurance model [16] or, from basic science, deriving tailored treatments and generating evidence by pain researchers is needed. Incorporating spontaneous recovery from initial pain, saluto-genetic models and patient preferences for specific interventions [17] should be taken into future considerations.

### **Tertiary Prevention of Pain: Reducing Disability, Work Loss, Negative Emotionality and Social Isolation for Sufferers with Chronic Pain**

Tertiary prevention aims to reduce secondary effects of those suffering from chronic pain. Models such as the fear avoidance model [18, 19] or the avoidance and endurance model [16, 20-23] incorporate the functional, psychological and social impairment in leading a satisfying life by the affected individuals. The biopsychosocial model of pain [24-26] led to the development of biopsychosocial (synonymously multidisciplinary, interdisciplinary, multi component) treatment approaches. A corresponding definition was devised by an IASP Task Force in 2017 and published at IASP homepage (<https://www.iasp-pain.org/Education/Content.aspx?ItemNumber=1698>).

Mayer and Gatchel [27] introduced the biopsychosocial treatment approach in the 1980s. Its main aim is the restoration of physical, psychological and social functioning, involving a core team of multidisciplinary healthcare professionals (physicians, physiotherapists, occupational therapists, psychologists and nurses) working in an integrated team setting [27].

Interdisciplinary treatment has been acknowledged as an appropriate answer to the comprehensive suffering of patients with chronic pain worldwide [28], but evidence is controversially discussed [29]. The heterogeneity of involved professions, providing interventions and the treatment composition, dose and duration of treatment, and the outcome assessment tools in clinical trials hampers considerably comparative effectiveness research and valid meta analyses [30]. Conceptual frameworks and mechanism based treatment designs, harmonized outcome assessment (considering comprehensive multidimensional outcomes [31]), carefully reported studies and carefully performed systematic research is needed to distinguish beneficial treatment from non-beneficial and, finally, to identify best treatment to specific groups of patients regarding their characteristics [32].

## REFERENCES

- [1] Loisel P. Developing a new paradigm: Work disability prevention. *Occupational Health Southern Africa* 2009;15(2):56-60.
- [2] Linton SJ. Chronic pain: the case for prevention. *Behaviour research and therapy* 1987;25(4):313-317.
- [3] Gewandter JS, Dworkin RH, Turk DC, Farrar JT, Fillingim RB, Gilron I, Markman JD, Oaklander AL, Polydefkis MJ, Raja SN, Robinson JP, Woolf CJ, Ziegler D, Ashburn MA, Burke LB, Cowan P, George SZ, Goli V, Graff OX, Iyengar S, Jay GW, Katz J, Kehlet H, Kitt RA, Kopecky EA, Malamut R, McDermott MP, Palmer P, Rappaport BA, Rauschkolb C, Steigerwal I, Tobias J, Walco GA. Research design considerations for chronic pain prevention clinical trials: IMMPACT recommendations. *Pain* 2015;156(7):1184-1197.
- [4] Fordyce WE, Brockway JA, Bergman JA, Spengler D. Acute back pain: a control-group comparison of behavioral vs traditional management methods. *Journal of Behavioral Medicine* 1986;9(2):127-140.
- [5] Meyer C, Denis CM, Berquin AD. Secondary prevention of chronic musculoskeletal pain: A systematic review of clinical trials. *Ann Phys Rehabil Med* 2018;61(5):323-338.
- [6] Winkelmann C and Schreiber T. Using 'White Flags' to categorize socio-cultural aspects in chronic pain. *European Journal of Public Health* 2019;29(Supplement\_4):ckz186-196.
- [7] Sowah D, Boyko R, Antle D, Miller L, Zakhary M, Straube S. Occupational interventions for the prevention of back pain: Overview of systematic reviews. *J Safety Res* 2018;66:39-59.
- [8] Sarmanova A, Fernandes GS, Richardson H, Valdes AM, Walsh DA, Zhang W, Doherty M. Contribution of central and peripheral risk factors to prevalence, incidence and progression of knee pain: a community-based cohort study. *Osteoarthritis Cartilage* 2018;26(11):1461-1473.
- [9] Melloh M, Elfering, Egli Presland C, Röder C, Hendrick P, Darlow B, Theis J-C. Predicting the transition from acute to persistent low back pain. *Occup Med (Lond)* 2011;61(2):127-31.
- [10] Johannsen M, Frederiksen Y, Jensen AB, Zacharie R. Psychosocial predictors of posttreatment pain after nonmetastatic breast cancer treatment: a systematic review and meta-analysis of prospective studies. *J Pain Res* 2018;11:23-36.
- [11] Kehlet H, Jensen TS, Woolf CJ, Persistent postsurgical pain: risk factors and prevention. *The Lancet* 2006;367(9522):1618-1625.
- [12] Reinpold W. Risk factors of chronic pain after inguinal hernia repair: a systematic review. *Innov Surg Sci* 2017;2(2):61-68.
- [13] Hill JC, Dunn KM, Main CJ, Hay EM. Subgrouping low back pain: a comparison of the STarT Back Tool with the Orebro Musculoskeletal Pain Screening Questionnaire. *Eur J Pain* 2010;14(1):83-89.
- [14] Hill JC, Dunn KM, Lewis M, Mullis R, Main CJ, Foster NE, Hay EM. A primary care back pain screening tool: identifying patient subgroups for initial treatment. *Arthritis Rheum* 2008;50(5):632-641.
- [15] Darlow B, Fullen BM, Dean S, Hurley DA, Baxter GD, Dowell A. The association between health care professional attitudes and beliefs and the attitudes and beliefs, clinical management, and outcomes of patients with low back pain: a systematic review. *Eur J Pain* 2012;16(1):3-17.
- [16] Hasenbring MI, Chehadi O, Titze C, Kreddig N. Fear and anxiety in the transition from acute to chronic pain: there is evidence for endurance besides avoidance. *Pain manage* 2014;4(5):363-374.
- [17] Aboagye E, Hagber J, Axén I, Kwak L, Lohela-Karlsson, M, Skillgate E, Dahlgren G, Jensen I. Individual preferences for physical exercise as secondary prevention for non-specific low back pain: A discrete choice experiment. *PLoS One* 2017;12(12):e0187709.
- [18] Crombez G, Eccleston C, Van Damme S, Vlaeyen JWS, Karoly P. Fear-avoidance model of chronic pain: the next generation. *Clin J Pain* 2012;28(6):475-483.
- [19] Vlaeyen JW and Linton SJ. Fear-avoidance model of chronic musculoskeletal pain: 12 years on. *Pain* 2012; 153(6):1144-1147.
- [20] Hasenbring MI, Hallner D, Klasen B, Streitlein-Böhme I, Willburger R, Rusche H. Pain-related avoidance versus endurance in primary care patients with subacute back pain: psychological characteristics and outcome at a 6-month follow-up. *Pain* 2012;153(1):211-217.
- [21] Hasenbring MI and Verbunt JA. Fear-avoidance and endurance-related responses to pain: new models of behavior and their consequences for clinical practice. *Clin J Pain* 2010;26(9):747-753.
- [22] Plaas H, Sudhaus S, Willburger R, Hasenbring MI. Physical activity and low back pain: the role of subgroups based on the avoidance-endurance model. *Disabil Rehabil* 2014;36(9):749-755.
- [23] Sudhaus S, Held S, Schoofs D, Bültmann J, Dück, I, Wolf OT, Hasenbring MI. Associations between fear-avoidance and endurance responses to pain and salivary cortisol in the context of experimental pain induction. *Psychoneuroendocrinology* 2015; 52:195-199.
- [24] Engel GL. The need for a new medical model: a challenge for biomedicine. *Science* 1977; 196(4286):129-136.
- [25] Engel GL. The clinical application of the biopsychosocial model. *J Med Philos* 1981;6(2):101-124.
- [26] Williams ACC and Craig KD. Updating the definition of pain. *Pain* 2016;157(11):2420-2423.

- [27] Mayer TG and Gatchel RJ. Functional restoration in spinal disorders: The Sports Medicine Approach. Philadelphia: Lea and Febiger; 1988.
- [28] Schatman M. Interdisciplinary chronic pain management: international perspectives. Pain: Clinical Updates 2012;20(7): 1-5.
- [29] Dragioti E, Evangelou E, Larsson B, Gerdle B. Effectiveness of multidisciplinary programmes for clinical pain conditions: An umbrella review. J Rehabil Med 2018;50(9):779-791.
- [30] Kaiser U, Treede R-D, Sabatowski R. Multimodal pain therapy in chronic noncancer pain—gold standard or need for further clarification? Pain 2017;158(10):1853-1859.
- [31] Deckert S, Kaiser U, Kopkow C, Trautmann F, Sabatowski R, Schmitt J. A systematic review of the outcomes reported in multimodal pain therapy for chronic pain. Eur J Pain 2016;20(1):51-63.
- [32] Turk DC. Chronic pain and whiplash associated disorders: rehabilitation and secondary prevention. Pain Res Manag 2003;8(1):40-3.

## AUTHOR

Dr. Ulrike Kaiser  
Lead. Clinical psychologist  
Comprehensive Pain Center  
University Hospital Carl Gustav Carus Dresden  
Dresden, Germany

## REVIEWERS

Brona M. Fullen, PhD  
Associate Professor  
UCD School of Public Health  
Physiotherapy and Sports Science  
Dublin, Ireland

Professor Esther Pogatzki-Zahn, MD, PhD  
Department of Anesthesiology, Intensive Care and Pain Medicine  
University Hospital Muenster  
Muenster, Germany