



## PATHOPHYSIOLOGICAL ASSESSMENT OF NON-SPECIFIC BACK PAIN



### BACK PAIN CAN HAVE A CLEAR UNDERLYING PATHOLOGY

Various clearly defined pathologies can be associated with back pain, including autoimmune disorders, spinal infections, or osteoporotic fractures. These pathologies have relatively clear signs and symptoms and well-defined diagnostic work-ups that ideally lead to a definite diagnosis and targeted treatment.

### PATHOLOGY IS UNCLEAR IN ~90% OF BACK PAIN PATIENTS

Nevertheless, back pain in at least 90% of patients cannot be attributed to any specific pathology [6]. As a consequence, diagnostic labels are descriptive in nature, e.g. 'Low back pain, unspecified' (ME.84.2Z in the ICD-11). In many research studies, the description is equally vague (e.g. 'non-specific low back pain').



### THIS LARGE PROPORTION IS UNLIKELY A HOMOGENOUS GROUP

It is highly unlikely that this large category of patients constitutes a similarly unified category as e.g., the autoimmune disorder ankylosing spondylitis. The important question becomes which pathophysiological mechanisms contribute to back pain in this category and how such mechanisms can be identified when assessing individual patients.



### PATIENT ASSESSMENT ALLOWS BROAD PATHOPHYSIOLOGICAL CATEGORIZATION

Not all mechanisms are currently known nor can they necessarily be assessed in humans. Nevertheless, broad pathophysiological categories can, at least partly, already be inferred at this point from patient assessment.

Like anywhere in the body, pain in the back can be ascribed to three fundamental pathophysiological categories: nociceptive, neuropathic and nociplastic pain [4], which are not mutually exclusive.



ASSESSMENTS



nociceptive



neuropathic



nociplastic

### ASSESSMENT SUMMARY

### ROUTINE CLINICAL ASSESSMENTS and ADDITIONAL OPTIONS to better understand pathophysiology (not exhaustive)

History taking/  
clinical exam



Pain drawings



Questionnaires



Laboratory tests



Imaging, e.g.  
Magnetic Resonance Imaging  
(MRI)



Quantitative Sensory Testing  
(QST)  
[8]



# PATIENT EXAMPLES

Let's consider three patients: Alex, Billy, and Sam. They are between 45 and 55 years of age, have had low back pain for six months (i.e. they have chronic pain per definition [10]) and they have recently received MR imaging of the lumbar spine (albeit this might not be in line with recommendations [2,5]). For all three, the MRI shows mild disc degeneration and mild facet joint degeneration at L3/4 and L4/5 without nerve root compression or Modic changes. The GP now sends the three patients to your office for further assessment. How will you find out to which pathophysiological category their pain predominantly belongs?


## ALEX

Alex complains about **persistent, burning pain** in the lower lumbar region extending paravertebrally and into the buttock in a **diffuse pattern**. No radiation to the lower limbs. **Movement does not aggravate** the pain, but sometimes, Alex feels some **itching** when wearing tight belts or pants.

### Clinical examination:

- no sensory or motor deficits
- range of motion (ROM) somewhat reduced in lumbar flexion and extension
- **no clear mechanical pattern** of pain aggravation by movement
- local tenderness upon palpation in the midline at the level L4/L5


No clear mechanical pattern, but burning pain and itchiness... **neuropathic pain?**

 **Neuropathic pain questionnaire**, e.g. the Neuropathic Pain Symptom Inventory (NPSI) [1]

- Alex scores 63 points on the weighted NPSI [9]

**QST**

- **QST @back**: increased mechanical detection threshold, reduced pressure pain threshold, mechanical allodynia and hyperalgesia. No alterations in thermal detection or thermal pain thresholds, relative to normative data [7]
- **QST @hand** (control area): all tests are normal [8]

 Evidence for a **neuropathic** component

Classical signs of central sensitization (mechanical allodynia and hyperalgesia in a secondary zone, i.e., the skin) and of nerve damage (increased mechanical detection threshold). Nevertheless, neuropathic pain cannot be definitely diagnosed because a lesion or disease of the somatosensory system is not confirmed in Alex [3]. Also, the signs and symptoms cannot be ascribed to a specific nerve territory or dermatome.


## BILLY

Billy suffers from **episodic, right-sided paraspinal** and right buttock pain, **aggravated by movement** and **standing** for a long time and **relieved by sitting and walking**. Billy denies pain at rest but after intense physical activity s/he feels **stiff** the next morning for **20-30 minutes**.

### Clinical examination:

- symptoms can be **provoked with extension and rotation** of the lumbar spine
- local tenderness over L4/L5 facet joints and musculature

Symptoms provoked by segmental movements... **nociceptive pain?**

 **Detailed clinical assessment** according to Vining and colleagues [11] to confirm the impression and identify the likely nociceptive source

- history taking and clinical exam (3 or more of: > age 50, relief by walking, relief by sitting, paraspinal onset, positive extension-rotation test [11]) point to the facet joint as the most likely source of pain

 Evidence for a **nociceptive** component

No further assessments are needed because Billy most likely suffers from nociceptive pain. It is possible that there is an inflammatory component associated with the degenerative changes that result in the movement-associated allodynia. However, there is no indication to suspect systemic inflammation.

## SAM

Sam reports **pain fluctuating in intensity and location, sometimes extending** to the **buttocks** and left or right **posterior thigh**. The pain gets **worse with movement**, but Sam also experiences **pain sometimes at rest**, sometimes waking him/her up at night.

### Clinical examination:

- **diffuse tenderness** over lumbar spinous processes and paraspinal muscles
- lumbar **movement is painful** in the end-range in **all directions** and ROM is somewhat restricted
- no sign of sensory or motor impairment

Pain at rest, waking up at night... **inflammatory component?**

 **Blood test**

- negative for inflammatory markers

Pain extending in the buttock and legs... **neuropathic component?**

 **NPSI** weighted score is 48, i.e. inconclusive [9]

**QST**

- **QST @painful sites** (back and leg): reduced pain thresholds for all modalities, increased pain sensitivity and normal detection thresholds. No dynamic mechanical allodynia [7,8]
- **QST @hand**: similar pattern of hypersensitivities, but to a lesser extent [8]

 Evidence for a **nociplastic** component

There is no clear evidence for an inflammatory or neuropathic component. A potential nociceptive component remains unclear (because of the movement-related pain), but the exam according to Vining and colleagues [11] does not allow a clear classification. The widespread hypersensitivity (at the hand in addition to the back and leg), the increased spatial pain extent and potentially the fluctuating character point towards nociplastic pain.

## OVERALL CONCLUSION

Alex, Billy and Sam represent relatively clear examples of pathophysiological pain categories. In reality, an individual's pain might of course arise from a mix of different pathophysiologicals. In addition, each category ought to be comprised of different mechanisms, which themselves are again the product of different sub-mechanisms. It is currently unclear in how much detail pathophysiology has to be understood to be most relevant for treatment; this will also depend at which mechanism treatment is targeted. Nevertheless, contributions of different pathophysiologicals to an individual's 'non-specific' back pain can be identified with assessment methods that are already available. The ultimate goal should be to get rid of the unfortunate diagnostic label of 'non-specific' back pain and, by better understanding of pathophysiological mechanisms, to develop and promote more targeted treatment in the future.



HISTORY TAKING & CLINICAL EXAM



ADDITIONAL ASSESSMENTS



CATEGORIZATION

## REFERENCES

- [1] Bouhassira D, Attal N, Fermanian J, Alchaar H, Gautron M, Masquelier E, Rostaing S, Lanteri-Minet M, Collin E, Grisart J, Boureau F. Development and validation of the Neuropathic Pain Symptom Inventory. *Pain* 2004;108(3):248-57.
- [2] Bussieres AE, Stewart G, Al-Zoubi F, Decina P, Descarreaux M, Haskett D, Hincapie C, Page I, Passmore S, Srbely J, Stupar M, Weisberg J, Ornelas J. Spinal Manipulative Therapy and Other Conservative Treatments for Low Back Pain: A Guideline From the Canadian Chiropractic Guideline Initiative. *J Manipulative Physiol Ther* 2018;41(4):265-93.
- [3] Finnerup NB, Haroutounian S, Kamerman P, Baron R, Bennett DL, Bouhassira D, Cruccu G, Freeman R, Hansson P, Nurmikko T, Raja SN, Rice AS, Serra J, Smith BH, Treede RD, Jensen TS. Neuropathic pain: an updated grading system for research and clinical practice. *Pain* 2016;157(8):1599-606.
- [4] International Association for the Study of Pain. Task Force on Taxonomy. IASP Terminology Updated from "Part III: Pain Terms, A Current List with Definitions and Notes on Usage" (pp 209-214), *Classification of Chronic Pain, 2nd Edition*. Seattle: IASP, 2017. p. IASP
- [5] Jenkins HJ, Downie AS, Maher CG, Moloney NA, Magnussen JS, Hancock MJ. Imaging for low back pain: is clinical use consistent with guidelines? A systematic review and meta-analysis. *Spine J* 2018;18(12):2266-77.
- [6] Maher C, Underwood M, Buchbinder R. Non-specific low back pain. *Lancet* 2017;389(10070):736-47.
- [7] Pfau DB, Krumova EK, Treede RD, Baron R, Toelle T, Birklein F, Eich W, Geber C, Gerhardt A, Weiss T, Magerl W, Maier C. Quantitative sensory testing in the German Research Network on Neuropathic Pain (DFNS): reference data for the trunk and application in patients with chronic postherpetic neuralgia. *Pain* 2014;155(5):1002-15.
- [8] Rolke R, Baron R, Maier C, Tolle TR, Treede RD, Beyer A, Binder A, Birbaumer N, Birklein F, Botefur IC, Braune S, Flor H, Huge V, Klug R, Landwehrmeyer GB, Magerl W, Maihofner C, Rolko C, Schaub C, Scherens A, Sprenger T, Valet M, Wasserka B. Quantitative sensory testing in the German Research Network on Neuropathic Pain (DFNS): standardized protocol and reference values. *Pain* 2006;123(3):231-43.
- [9] Sommer C, Richter H, Rogausch JP, Frettlöh J, Lungenhausen M, Maier C. A modified score to identify and discriminate neuropathic pain: a study on the German version of the Neuropathic Pain Symptom Inventory (NPSI). *BMC Neurol* 2011;11:104.
- [10] Treede RD, Rief W, Barke A, Aziz Q, Bennett MI, Benoliel R, Cohen M, Evers S, Finnerup NB, First MB, Giamberardino MA, Kaasa S, Kosek E, Lavand'homme P, Nicholas M, Perrot S, Scholz J, Schug S, Smith BH, Svensson P, Vlaeyen JW, Wang SJ. A classification of chronic pain for ICD-11. *Pain* 2015;156(6):1003-7.
- [11] Vining RD, Minkalis AL, Shannon ZK, Twist EJ. Development of an Evidence-Based Practical Diagnostic Checklist and Corresponding Clinical Exam for Low Back Pain. *J Manipulative Physiol Ther* 2019.

## AUTHORS

### Petra Schweinhardt, MD, PhD \*

Integrative Spinal Research  
Department of Chiropractic Medicine  
Balgrist University Hospital  
University of Zurich  
Zurich, Switzerland

University of Zurich  
Zurich, Switzerland

### Luana Nyiroe, DCM

Integrative Spinal Research  
Department of Chiropractic Medicine  
Balgrist University Hospital  
University of Zurich  
Zurich, Switzerland

University of Zurich  
Zurich, Switzerland

### Mirjam Baechler, DC, MMEd

Integrative Spinal Research  
Department of Chiropractic Medicine  
Balgrist University Hospital  
University of Zurich  
Zurich, Switzerland

University of Zurich  
Zurich, Switzerland

### Laura Sirucek, MSc

Integrative Spinal Research  
Department of Chiropractic Medicine  
Balgrist University Hospital  
University of Zurich  
Zurich, Switzerland

University of Zurich  
Zurich, Switzerland

### Susanne Becker, PhD

Integrative Spinal Research  
Department of Chiropractic Medicine  
Balgrist University Hospital  
University of Zurich  
Zurich, Switzerland

University of Zurich  
Zurich, Switzerland

Department of Cognitive and Clinical Neuroscience  
Medical Faculty Mannheim  
Central Institute of Mental Health  
Heidelberg University  
Mannheim, Germany

\*corresponding author: [petra.schweinhardt@balgrist.ch](mailto:petra.schweinhardt@balgrist.ch)

## REVIEWERS

### Owen D Williamson, FRCSC Pain Medicine

Adjunct Professor  
School of Interactive Arts and Technology  
Simon Fraser University  
Surrey, BC, Canada

### Thomas Graven-Nielsen, DMSc, PhD

Center for Neuroplasticity and Pain (CNAP)  
Aalborg University, Denmark

## ACKNOWLEDGMENTS

This work is supported by a Clinical Research Priority Program of the University of Zurich (CRPP Pain).